Local Photon and Graviton Mass and its Consequences

M. Tajmar*

ARC Seibersdorf research GmbH, A-2444 Seibersdorf, Austria

C. J. de Matos[†]

ESA-HQ, European Space Agency, 8-10 rue Mario Nikis, 75015 Paris, France

Abstract

We show that the presently accepted value of the cosmological constant and a correspondingly small graviton mass leads to considerable gravitomagnetic fields around rotating mass densities, which are not observed. The solution to the problem is found by a graviton mass which depends on the local mass density to ensure the principle of equivalence. This solution, derived from Einstein-Proca equations, has important consequences such as the correct prediction of the dark energy density in the universe solving the "cosmic coincidence" problem, a prediction of the Higgs mass in line with present estimates and a correction term for the Cooper-pair mass anomaly reported by Tate among many others. Perhaps the most interesting results are that the vacuum energy density is then proportional to the energy density of matter in our universe; and that coherent matter can be used to engineer the vacuum. Similar results were obtained for the photon mass which is then proportional to the charge density in matter. For the case of coherent matter the predicted effects have been experimentally observed by the authors.

PACS: 98.80.Bp, 98.80.Es

Keywords: Graviton, Cosmological Constant, Dark Energy, Coincidence Problem, Higgs Mass, Cooper-Pair Mass Anomaly.

^{*} Head of Space Propulsion, Phone: +43-50550-3142, Fax: +43-50550-3366, E-mail: martin.tajmar@arcs.ac.at

[†] General Studies Officer, Phone: +33-1.53.69.74.98, Fax: +33-1.53.69.76.51, E-mail: clovis.de.matos@esa.int

1. Introduction

It is well known that the mass of the photon and graviton in vacuum must be non-zero. The first limit is given by Heisenberg's uncertainty principle¹ and the second by the measurement of the cosmological constant in our universe²-⁴. Both Heisenberg's limit and the cosmological constant lead respectively to a photon and a graviton mass of about 10⁻⁶⁹ kg. This is obviously very small and it is therefore believed that it has negligible consequences. We will show that this is actually not the case and leads to fundamental new insights both for classical and quantum matter. An immediate consequence is that Maxwell equations transform into Proca type equations leading us to the conclusion that for the electromagnetic and gravitational interaction gauge invariance only applies to a certain approximation in free space. The consequence of this result will be discussed in the following. Especially the new treatments required for the graviton have far reaching consequences that can be experimentally assessed.

2. Proca Equations and the Photon Mass

As the photon's mass is non-zero, the usual Maxwell equations for electromagnetism transform into the well known Proca equations with additional terms due to the finite Photon wavelength

$$div \vec{E} = \frac{\rho}{\varepsilon_0} - \left(\frac{m_{\gamma}c}{\hbar}\right)^2 \cdot \varphi$$

$$div \vec{B} = 0$$

$$rot \vec{E} = -\frac{\partial \vec{B}}{\partial t}$$

$$rot \vec{B} = \mu_0 \rho \vec{v} + \frac{1}{c^2} \frac{\partial \vec{E}}{\partial t} - \left(\frac{m_{\gamma}c}{\hbar}\right)^2 \cdot \vec{A}$$
(1)

The usual effect attributed to a finite Photon mass is its consequence on the strength of electromagnetic forces over distances. However, taking the curl of the 4th equation reveals another feature, which was first assessed by the authors in the framework of superconductivity⁵:

$$B = B_0 \cdot e^{-\frac{x}{\lambda_{\gamma}}} + 2\omega\rho\mu_0\lambda_{\gamma}^2 , \qquad (2)$$

The first part of Equ. (2) is the Yukawa-type exponential decay of the magnetic field, and the second part shows that a magnetic field will be generated due to the rotation of a charge density ρ . In quantum field theory, superconductivity is explained via a large photon mass as a consequence of gauge symmetry breaking and the Higgs mechanism. The photon wavelength is then interpreted as the London penetration depth and leads to a Photon mass about 1/1000 of the electron mass. This then leads to

$$B = B_0 \cdot e^{-\frac{x}{\lambda_L}} - 2\frac{m^*}{e^*}\omega , \qquad (3)$$

where the first term is called the Meissner-Ochsenfeld effect (shielding of electromagnetic fields entering the superconductor) and the second term is known as the London moment (minus sign is due to the fact that the Cooper-pairs lag behind the rotation) with m^* and e^* as the Cooper-pair's mass and charge. The magnetic field produced by a rotating superconductor as a consequence of its large photon mass was experimentally measured outside of the quantum condensate where the photon mass is believed to be close to zero^{8,9}.

Knowing the effect from a large photon mass in superconductivity, what is the effect of a non-zero photon mass in normal matter? Contrary to the case of superconductivity where the photon mass in comparable to the mass on an electron, in normal matter, the photon mass is believed to be close to zero. The problem of the Proca equations is easily shown: The "Meissner" part becomes important only for large photon masses (which is not the case in normal matter) – but the "London Moment" part becomes important for very small photon masses. By taking the presently accepted experimental limit on the photon mass¹ (m_{γ} < 10^{-52} kg), we can write the second part as

$$B > \omega \rho \cdot 2.8 \times 10^{13} . \tag{4}$$

This would mean that a charge density ρ rotating at an angular velocity ω should produce huge magnetic fields. Obviously, this is not the case leading to a paradox for normal matter. Therefore the value of the photon mass for matter containing a charge density must be different from the one in free space.

In fact, there is only one possible choice for the photon mass. Let's consider the case of a single electron moving in a magnetic field. It will then perform a precession movement according to Larmor's theorem ω =-(q/2m).B. We can then solve Equ. (2) for the photon mass expressing it as

$$\frac{1}{\lambda_{\gamma}^{2}} = \left(\frac{m_{\gamma}c^{2}}{\hbar}\right)^{2} = -\frac{q}{m} \cdot \mu_{0}\rho . \tag{5}$$

We find that the photon mass inside normal matter must be defined over the charge density and the charge-to-mass ratio observed. Note that the photon mass is then (due to the negative sign in the Larmor theorem) always a complex value independent of the sign of charge. This is a surprising result and will need further investigation which is under preparation by the authors. Especially as the photon mass inside a superconductor due to the Higgs mechanism has a real value. As a hypothesis, we can postulate that the complex mass will change into a real value by passing from normal to coherent matter. That would actually introduce a sign change in the London moment and will not require Becker's argument¹⁷ any more to get the right sign as observed in experiments.

Only for neutral matter or vacuum, the photon mass can be therefore given by the limit obtained via Heisenberg's uncertainty principle. As nearly all matter in the universe can be considered neutral, this consequence may be of minor importance. However, in case of gravity and the graviton, the consequences are far reaching as matter is never neutral in a gravitational sense.

3. Proca Equations and Graviphoton Mass

In the weak field approximation, gravity can be written similar to a Maxwellian structure forming the so-called Einstein-Maxwell equations. The quantization of the Maxwellian theory of gravity would lead to a spin one boson as a mediator of gravitoelectromagnetic fields. This represents a major problem with respect to the theory of general relativity, which only predicts quadrupolar gravitational waves associated with spin two gravitons. This is the reason why the linear approximation of Einstein field equations is taken as being only an approximation to the complete theory, which cannot be used to investigate radiative processes. Recent experimental results on the gravitomagnetic London moment⁶ tend to demonstrate that gravitational dipolar type radiation associated with the Einstein-Maxwell equations is real. This implies that Maxwellian gravity is not only an approximation to the complete theory, but may indeed reveal a new aspect of gravitational phenomena associated with a vectorial spin 1 gravitational boson, which we might call the graviphoton. As an example, a fully relativistic modified theory of gravity called scalar vector tensor gravity would duly take into account this new side of gravity. In the following we will therefore use the term graviphoton for studying the Proca type character of gravity and its consequences on cosmology, coherent matter and high energy particle physics.

Similar to the case of the photon, the graviphoton mass is not zero based on the measurement of the cosmological constant. The field equations for massive linearized gravity are given by¹¹:

$$\begin{aligned}
div \, \bar{g} &= -\frac{\rho_m}{\varepsilon_g} - \left(\frac{m_g c}{\hbar}\right)^2 \cdot \varphi_g \\
div \, \bar{B}_g &= 0 \\
rot \, \bar{g} &= -\frac{\partial \bar{B}_g}{\partial t} \\
rot \, \bar{B}_g &= -\mu_{0g} \, \rho_m \vec{v} + \frac{1}{c^2} \frac{\partial \bar{g}}{\partial t} - \left(\frac{m_g c}{\hbar}\right)^2 \cdot \vec{A}_g
\end{aligned}$$
(6)

where g is the gravitoelectric and B_g the gravitomagnetic field. The authors want to stress that Equs. (6) are intriguingly similar to the set of Proca equations obtained from the linearization of Einstein field equations with a cosmological constant¹². The recent measurement of the cosmological constant $\Lambda = (1.29 \pm 0.23) \times 10^{-52}$ m² by WMAP² can be linked to the graviton (graviphoton) mass by a recent result from Novello et all and others^{3,4}

$$m_g = \frac{\hbar}{c} \cdot \sqrt{\frac{2}{3}\Lambda} = 3 \times 10^{-69} \quad [kg].$$
 (7)

This result should not surprise the reader since as mentioned above the origin of Equs. (6) seem to be Einstein field equations with a cosmological constant. There is a debate going on in the literature if the graviton (graviphoton) is a real or complex number. We will see later that it has to be a real number which is also confirmed by our experimental results⁶. Applying again a curl on the 4th Proca equation leads to

$$B_{g} = B_{0,g} \cdot e^{-\frac{x}{\lambda_{g}}} - 2\omega \rho_{m} \mu_{0g} \lambda_{g}^{2} , \qquad (8)$$

where ρ_m is now the mass density and μ_{0g} the gravitomagnetic permeability (note the different sign with respect to Equ. (3)). Similar to electromagnetism, we obtain a Meissner and a London moment part for the gravitomagnetic field generated by matter⁵.

4. Consequences of Local Graviton Mass

Applying the graviton mass in Equ. (7) to Equ. (8) leads again to huge gravitomagnetic fields for rotating mass densities which are not observed. Our choice for the graviton mass was of course due to the measurement of the cosmological constant, a value determined from supernovae at the outer boundaries of the universe and anisotropies in the cosmic microwave background radiation². Therefore the result should be valid at least on the scale of the universe, but might not apply locally to ordinary matter in the laboratory.

Again we find the solution in the gravitational analog to the Larmor theorem. Locally, the principle of equivalence must be fulfilled for any type of matter. That means that local accelerations must be equivalent to gravitational fields and a body can not distinguish between being in a rotating reference frame or being subjected to a gravitomagnetic field ($B_{\rm g}=-2\omega$).

By choosing the graviton wavelength proportional to the local density of matter similar to Equ. (5),

$$\frac{1}{\lambda_g^2} = \left(\frac{m_g c}{\hbar}\right)^2 = \mu_{0g} \rho_m , \qquad (9)$$

we find the solution. Note that due to the different signs in the Einstein-Proca equations, we find that the graviton mass is a real number to comply with the Larmor theorem in normal matter. Inserting Equ. (9) into our Proca equations for gravity in Equ. (6), we find by performing another grad operator on the 1st equation and another curl on the 4th equation

$$g = -a, \quad B_g = -2\omega \quad , \tag{10}$$

which is nothing else as the formulation of the equivalence principle and the gravitomagnetic Larmor theorem¹⁰. The graviton mass in Equ. (9) therefore describes the inertial properties of matter in accelerated reference frames. This is a very fundamental result and new insight into the foundations of mechanics. It can be also interpreted as a form of Mach's principle.

A similar graviton mass (up to a factor 3/16) was already found by Argyris¹¹ by solving Einstein's equations in the conformally flat case. He linked it with the average mass of the universe. However, as we have shown, this result is valid locally for all matter. As we will show now, this graviton mass has far reaching consequences in determining dark energy as well as deviations from the classical gravitomagnetic Larmor theorem (2nd equation in Equ. (10)).

It is interesting to note that if we take the case of no local sources ($\rho_{\scriptscriptstyle m}=0$), the graviphoton mass will be zero, and we will find, by solving the weak field equations in

the transverse gauge, the "classical" freely propagating degrees of freedom of gravitational waves associated with a massless spin 2 graviton¹³. However, in the case of local sources, a spin-1 graviphoton will appear.

4.1 Application to Normal Matter

Following Equ. (7) we can assess the consequences of the local graviton mass for our present models of the universe. The cosmological constant for Einstein's equation then is the given by

$$\Lambda = \frac{3}{2} \frac{1}{\lambda_g^2} = \frac{3}{2} \mu_{0g} \rho_m . \tag{11}$$

The average density of the universe² is $\rho_m \approx 10^{-26}$ kg.m⁻³. This gives a graviton mass of $m_g \approx 3.2 \times 10^{-69}$ kg and a cosmological constant, using Equ. (11), of $\Lambda \approx 1.3 \times 10^{-52}$ m⁻². Those values are exactly within present experimental observations! How large is the local graviton mass? In a piece of iron for example, the absolute value of the graviton mass would be 2.8×10^{-54} kg, which is still undetectable small. **Table 1** lists the cosmological constant based on our calculations for some cases^{*}.

We can also use Equ. (11) to express the vacuum energy density as

$$\rho_V = \frac{3}{4} \rho_m c^2 \ . \tag{12}$$

So in fact, the vacuum energy density is equal (up to the numerical factor of 0.75) to the energy density of matter ($\rho_E = \rho_m c^2$). This is indeed an intriguing result and gives a totally new perspective to the energy of the vacuum – being defined as function of the local density of matter.

These results can be used to actually solve the cosmological coincidence problem (observations showed that dark energy makes up about 73% of the energy present in the universe). In a flat universe, which is presently observed, the average density of the universe can be expressed by

$$\rho_m = \frac{3H^2}{8\pi G} \ . \tag{13}$$

The dark energy density is expressed by

$$\Omega_{\Lambda} = \frac{\Lambda c^2}{3H^2} \ . \tag{14}$$

By combining Equs. (12)-(14) and using $\mu_{0g} = \frac{4\pi G}{c^2}$ we get

$$\Omega_{\Lambda} = \frac{3}{4} = 0.75 . \tag{15}$$

This fits remarkably well to present observations² of Ω_{Λ} =0.73±0.04, it is obviously the same factor that is in front of the mass energy density in Equ. (12). So the answer to

^{*} Note that for the case of the solar system, the orbit of pluto is 5 orders of magnitude less than the wavelength of

the cosmological coincidence is that it is a natural consequence of the fact that the graviton mass is a function of the matter density and a flat universe. We can use this analytical result also to express a direct relationship between the Hubble and cosmological constant

$$H^2 = \frac{4}{9}\Lambda c^2 \ . \tag{16}$$

Of course a graviton mass has other implications. We will assess here just some of the many consequences. An obvious one is the dispersion of gravitational waves depending on the local mass density. A graviton mass leads to a frequency dependence on the propagation of gravity in free space. This has already been assessed by others^{1,11}. Using our Equ. (9), we can write the group velocity of gravity as

$$v_g = c \cdot \sqrt{1 - 4\pi G \frac{\rho_m}{\omega^2}} \quad . \tag{17}$$

For low frequencies and high densities, this factor can become important. Another consequence is that a non-zero cosmological constant has also an influence on the Schwarzschild solution for black holes. The Schwarzschild metric in de-Sitter spacetime can be written as 14

the graviphoton in the solar system.

$$d\tau^{2} = \left(1 - \frac{2MG}{c^{2}r} - \frac{\Lambda r^{2}}{3}\right)dt^{2} - \left(1 - \frac{2MG}{c^{2}r} - \frac{\Lambda r^{2}}{3}\right)^{-1}dr^{2} - r^{2}\left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right). \tag{18}$$

Using the expression for the cosmological constant in Equ. (11), we can therefore express the Schwarzschild radius using the metric in Equ. (18) as

$$r_{\rm S} = \frac{7MG}{2c^2} \ . \tag{19}$$

This deviates from the classical result by a factor of 1.75, which is small. However, there is an important difference with respect to previous work on this subject. Previously it was thought that the gravitational horizon $L=\Lambda^{-0.5}$ is always much larger than the Schwarzschild radius, because a cosmological constant of 1.29x10⁻⁵² m² was assumed. In our case, the cosmological constant of a black hole is given by

$$\Lambda_S = \frac{36c^4}{343M^2G^2} \ . \tag{20}$$

This gives a gravitational horizon of

$$L_{S} = \frac{1}{\sqrt{\Lambda_{S}}} = \frac{\sqrt{343}}{6} \frac{MG}{c^{2}} , \qquad (21)$$

which is just a little bit smaller than the Schwarzschild radius of Equ. (19). This has the consequence, that the gravitational force inside the black hole should now decrease with a Yukawa type modification affecting gravitational and gravitomagnetic fields. The denser the black hole, the stronger will be the gravitational force decay.

This regulation mechanism of course can lead to a totally new fate for black holes different from present predictions. In fact, it has already been shown that a spin-1 field such as the graviphoton in Einstein-Proca equations prohibits black hole solutions¹⁵. That can be an important input to solve presently observed anomalies such as nested discs in rotating Keplerian rotation around supermassive black holes¹⁶.

4.2 Application to Coherent Quantum Matter

Perhaps that most important consequence of the local graviton mass is its relation to superconductivity. As we wrote already in the introduction of this paper, the application of Proca equations to superconductivity are well established. In a superconductor, we have now a ratio between matter being in normal and in a condensed (coherent) state. So we have two sets of Proca equations, one which deals with the overall mass and one with its condensated subset.

The coherent part of a given material (e.g. the Cooper-pair fluid) is also described by its own set of Proca equations similar to the ones in Equ. (6) but with one important difference: Instead of the ordinary mass density ρ_m we have to take the Cooper-pair mass density ρ_m^* . By taking the curl of the fourth Proca equation (See Ref 5), we arrive at

$$B_{g} = B_{0,g} \cdot e^{-\frac{x}{\lambda_{g}}} - 2\omega \rho_{m}^{*} \mu_{0g} \lambda_{g}^{2} , \qquad (22)$$

where we had to introduce Becker's argument¹⁷ that the Cooper-pairs are lagging behind the lattice so that the current is flowing in the opposite direction of ω . This gives the right sign with respect to our experimental observation⁶. A similar argument was introduced for the classical London moment as also here the sign change was observed accordingly. The first part (Meissner part) is not different from our previous assessment for normal matter, but the second part changes to

$$B_{g} = 2\omega \frac{\rho_{m}^{*}}{\rho_{m}} , \qquad (23)$$

due to the fact that the graviton mass depends on all matter in the material, not just the coherent part. That is why the densities do not cut any more and we arrive at an additional field for coherent matter in addition to the gravitational Larmor theorem in Equ. (10). We switched here from the real graviton (graviphoton) mass to a complex value similar to what we discussed for the photon. This gives again the right sign for the gravitomagnetic London moment as observed and avoids Becker's argument¹⁷.

Similarly, by taking the gradient of the first Proca equation in Equs (6), considering the case of an homogeneous gravitational field, and using the fact that the gradient of a density of energy is equal to a density of force, $\nabla(\rho_m^*c^2) = \rho_m^*\vec{a}$, we obtain:

$$\vec{g} = \vec{a}\mu_{0g}\rho_m^*\lambda_g^2 , \qquad (24)$$

which transforms using Equ.(9), into:

$$\vec{g} = -\frac{\rho_m^*}{\rho_m} \vec{a} , \qquad (25)$$

where \vec{a} is the total net acceleration to which the superconductor is submitted.

Comparing Equs (23) and (25) to Equs (10) we conclude that the presence of Cooper-pairs inside the superconductor leads to a deviation from the equivalence principle and from the classical gravitational Larmor theorem. A rigid reference frame mixed with non-coherent and coherent matter is not equivalent to a rigid reference frame made of normal matter alone, with respect to its inertial and gravitational properties. However, in the case of a Bose-Einstein condensate where we have only coherent matter, Equ. (25) transforms into the usual expression for normal matter and the equivalence principle is again conserved.

An important feature is that these fields (Equs. (23) and (25)) however, should be also present <u>outside</u> the superconductor, contrary to the classical inertial behaviour. The gravitomagnetic Larmor theorem for normal matter describes the inertial forces in an accelerated reference frame. This leads to so-called pseudo forces which are only present inside the material which is rotating. The difference to quantum materials is that in this case, the integral of the canonical momentum is quantised. Let's consider a superconducting ring. The integral of the full canonical momentum of the Cooper-pairs including gravitational fields is given by

$$\oint \vec{p}_s \cdot d\vec{l} = \oint \left(m^* \vec{v}_s + e^* \vec{A} + m^* \vec{A}_g \right) \cdot d\vec{l} = \frac{nh}{2} .$$
(26)

If the ring is thicker than the London penetration depth, then the integral can be set to zero. Solving for the case where the superconductor is at an angular velocity ω , we get

$$\vec{B} = -\frac{2m}{\rho} \cdot \vec{\omega} - \frac{m}{\rho} \cdot \vec{B}_g \,. \tag{27}$$

These magnetic and gravitomagnetic fields are also present inside the superconducting ring. The first part is the classical London moment, with its origin is due to the photon mass, and the second part is its analog gravitomagnetic London moment, which will produce an additional field overlapping the classical London moment. According to Equ. (23), depending on the superconductor's bulk and Cooper-pair density, the magnetic field should be higher than classically expected. Indeed, that has been measured without apparent solution throughout the literature.

The authors already conjectured such a field to explain a reported disagreement between the theoretical and experimental Cooper-pair mass 5,18,19 . Tate et al 8,9 used a sensitive London moment measurement to determine the Cooper-pair mass in Niobium. This mass was found to be larger ($m^*/2m_e = 1.000084(21)$) than the theoretically expected value ($m^*/2m_e = 0.999992$). As we pointed out earlier in our conjecture, this mass difference opens up the room for large gravitomagnetic fields following the quantised canonical momentum in Equ. (24). In order to correct Tate's result, we need a gravitomagnetic field of

$$B = 2 \frac{m_{\text{exp} erimental}^* - m_{\text{theoretical}}^*}{m_{\text{theo}}^*} \omega = 2 \frac{\Delta m^*}{m^*} \omega , \qquad (28)$$

where Δm^* is the difference between the experimental and theoretical Cooper-pair mass to find back the Cooper-pair mass predicted by quantum theory.

The local graviton mass now establishes the reason why such a gravitomagnetic field has to be there. Comparing Equs. (23) and (28), we can identify

$$\frac{\Delta m^*}{m^*} = \frac{\rho_m^*}{\rho_m} \quad . \tag{29}$$

Taking Tate's values ($\frac{\Delta m^*}{m^*} = 9.2 \times 10^{-5}$) and the Niobium bulk and Cooper-pair mass

density $(\frac{\rho_m^*}{\rho_m} = 3.95 \times 10^{-6})$, we see that these values are a factor of 23 away. One has to take into account that Tate's measurement is up to now the only precision

experiment and, even more important, other relativistic correction terms need to be

added to the theoretical Cooper-pair mass, which will make Δm^* smaller and Equ.

(29) match better.

This is a very important result as we have for the first time not only a conjecture to explain Tate's anomaly, but also a good reason why a rotating superconductor should produce a gravitomagnetic field which is larger than classical predictions from ordinary rotating matter. The reason is the local graviphoton mass. **Fig. 1** plots the predicted $\Delta m/m$ for various superconductors.

In parallel, the authors also attempted to measure the gravitomagnetic field from induced gravitational fields around a rotating superconductor which is reported in Ref. 6. First measurements show that this gravitomagnetic field indeed exists with a measurement in between our Cooper-pair density ratio and the one derived from Tate's measurements ($\frac{\Delta m^*}{m^*}$). This adds strong confidence in our theoretical approach and its consequences.

5. Mass of the Higgs Boson

As the last but not least consequence, we want to assess our results in the framework of the Higgs boson mass. According to the Standard Model (SM), the vacuum in which all particle interactions take place is not actually empty, but is instead filled with a condensate of Higgs particles. The quarks, leptons, and W and Z bosons continuously collide with these Higgs particles as they travel through the "vacuum". The Higgs condensate acts like molasses and slows down anything that interacts with it. The stronger the interactions between the particles and the Higgs condensate are, the heavier the particles become²⁰.

The Higgs mechanism is an essential part of the Standard Model. Without it the quarks and leptons - and also the W and Z bosons - would all be massless and the world as we know it, could not exist. However, the physics behind the Higgs mechanism is the least tested aspect of the Standard Model. Although we have much circumstantial evidence for the Higgs particle, given that fundamental particles have masses that are consistent with the Higgs mechanism and from indirect

measurements at CERN and Stanford (so-called precision electroweak data), Higgs particles have never been directly produced and observed in collider experiments²¹.

The standard model predicts 22 that the vacuum energy density is directly proportional to the square of the Higgs mass m_{H}

$$\left|\rho_{V}\right| = \frac{1}{2} \left(m_{H} v c\right)^{2} \tag{30}$$

where v is the Vacuum Expectation Value (VEV) of the Higgs mass. Delgado 23 found, that the VEV essentially measures the mean number density of Higgs particles n_H condensed in the zero-momentum state (vacuum),

$$v^2 = \frac{2n_H}{m_H} {(31)}$$

Assuming that a Cooper-pair condensate is a possible form of a Higgs condensate, we can equal the density of condensed Higgs particles n_H to the Cooper-pair density n_s in a superconductor. Doing Equ. (31) into Equ. (30) we obtain:

$$\left|\rho_{V}\right| \approx \frac{1}{2} n_{s} m_{H} c^{2} \tag{32}$$

This looks very similar to the vacuum energy relation that we found in Equ. (12). By equalling both, we can estimate the mass of the Higgs boson in function of the local density of mass

$$m_H \approx \frac{3}{2} \frac{\rho_m}{n_s} \tag{33}$$

Taking the example of Niobium (ρ_m =8570 kg.m⁻³, n_s =3.7x10²⁸ m⁻³), we estimate the Higgs mass as m_H =192 GeV. Measurements at CERN and Fermilab estimate the Higgs mass between 96 and 117 GeV/c² with an upper limit of 251 GeV/c². This is very close to our estimate above. This result leads us to consider coherent matter on the same physical foot as spacetime vacuum. Coherent matter would then be a form of vacuum.

It is interesting to note that from Equ.(9) and Equ.(33) we observe that the local density of mass determines respectively the local mass of the graviphoton as well as the local mass of the Higgs boson. On the other side by equalling both equations we deduce that the local Higgs boson mass is proportional to the square of the local graviphoton's mass.

$$m_H \propto m_g^2$$
 . (34)

This can be understood as being a fundamental bridge between linearized general relativity with a cosmological constant, and the standard model. The physical meaning of that bridge still needs to be explored.

Conclusion

We have shown that the presently accepted value of the cosmological constant and a correspondingly small graviton mass leads to huge gravitomagnetic fields around rotating mass densities, which are not observed. The solution to the problem is found by a graviton mass which depends on the local mass density leading to the correct inertial forces in rotating reference frames. That can be understood as a foundation of basic mechanics. This solution, derived from Einstein and Proca equations, has important consequences such as

- the vacuum energy is always given (up to a numerical factor of 0.75) by the energy density of matter independently of scale,
- the dark energy density in a flat universe is always 0.75, matching present observations and solving the so-called coincidence problem,
- for the case of normal matter, the Proca equations now lead to the gravitomagnetic Larmor theorem and not to unobserved huge gravitomagnetic fields,
- the prediction of a gravitomagnetic London moment (observed experimentally) in rotating superconductors, that can solve the Cooper-pair mass anomaly reported by Tate,
- the prediction of a Higgs mass in line with current observations,

among many others such as black holes and the dispersion of gravity as we briefly assessed. Similar results have also been outlined for the Photon mass. By obeying Larmor's theorem, we find that in classical matter the photon has a complex and the graviton(graviphoton) a real value. In coherent matter we suggest the hypothesis that

it is exactly the other way round, which solves the sign change problems associated to the classical and gravitomagnetic London moment. The results are very encouraging and shall stimulate the further development of the basic concept outlined in this paper: The utilization of coherent matter to engineer the vacuum.

References

¹Tu, L., Luo, J., Gillies, J.T., Rep. Prog. Phys. **68**, 77-130 (2005).

²Spergel, D.N., et al., *Astrophy. J. Suppl.*, **148**, 175 (2003).

³Novello, M., Neves, R.P., Class. Quantum Grav. **20**, L67-L73 (2003).

⁴Liao, L., gr-qc/0411122, 2004

⁵De Matos, C.J., Tajmar, M., *Physica C* **432**, 167-172 (2005).

⁶Tajmar, M., Plesescu, F., Marhold, K., de Matos, C.J., *Physica C (submitted)*

⁷Moffat, J.W., gr-qc//0506021

⁸Tate, J., Cabrera, B., Felch, S.B., Anderson, J.T., *Phys. Rev. Lett.* **62**(8), 845-848 (1989).

⁹Tate, J., Cabrera, B., Felch, S.B., Anderson, J.T., *Phys. Rev. B* **42**(13), 7885-7893 (1990).

¹⁰Mashhoon, B., *Phys. Lett. A*, **173**, 347-354 (1993).

¹¹Argyris, J., Ciubotariu, C., *Aust. J. Phys.*, **50**, 879-891 (1997).

¹²de Matos, C.J., Tajmar, M., to be published

¹³Carroll, S., Spacetime and Geometry: An Introduction to General Relativity (Addison Wesley; 1st edition, 2003), 293-300

¹⁴Neupane, I.P., gr-qc/9812096

¹⁵Obukhov, Y.N., Vlachynsky, E.J., *Ann. Phys.* **8**(6), 497 – 509 (1999).

¹⁶Bender, R., *The Astrophysical Journal* **631**, 280-300 (2005).

¹⁷Becker, R., Heller, G., Sauter, F., *Z. Physik* **85**, 772-787 (1933).

¹⁸Tajmar, M., de Matos, C.J., Gravitomagnetic Field of a Rotating Superconductor and of a Rotating Superfluid", *Physica C*, **385**(4), 2003, pp. 551-554

¹⁹Tajmar, M., de Matos, C.J., *Physica C* **420**(1-2), 56-60 (2005).

²⁰Rolnick, W.B., Fundamental Particles and their Interactions (Addison-Wesley Publishing Company, 1994)

²¹Renton, P., *Nature*, **428**, 141 (2004).

²²Delgado, V., hep-ph/9305249

²³Delgado, V., hep-ph/9403247

Location	Cosmological Constant [m²]
Sun	1.97x10 ⁻²³
Earth	7.68x10 ⁻²³
Solar System	3.14x10 ⁻³⁵
Milky Way	6.29x10 ⁻⁴⁸
Universe	1.29x10 ⁻⁵²

 Table 1
 Cosmological Constant Examples




Fig. 1 Predicted $\Delta m/m$ for various Superconductors